

Corlanor

United Arab Emirates · access guide

How to access Corlanor for chronic heart failure from the UAE: 2026 pathway via UAE cardiology and pharmacy supply | Reserve Meds

By Reserve Meds clinical & regulatory team. Last reviewed 2026-05-20.

The UAE has one of the deepest adult and paediatric cardiology service networks in the wider region. Cleveland Clinic Abu Dhabi Heart and Vascular Institute, Sheikh Shakhboud Medical City (SSMC), Sheikh Khalifa Medical City (SKMC), Burjeel Medical City cardiology, Mediclinic City Hospital Heart and Vascular Institute, American Hospital Dubai cardiology, and the Dubai Health Authority hospitals all run heart-failure clinics that move patients through the full guideline-directed medical therapy ladder: ACE inhibitor or ARB or ARNI (Entresto), evidence-based beta-blocker, mineralocorticoid receptor antagonist, SGLT2 inhibitor, and where the resting heart rate stays elevated on maximally tolerated beta-blocker, Corlanor (ivabradine, Amgen US license; Procoralan is the Servier-originated European brand of the same molecule, and may be the registered name in some Gulf markets). For a UAE-resident adult or paediatric patient with HFrEF and a resting sinus rate at or above 70 bpm despite optimised beta-blocker therapy, the operational question is no longer whether selective sinoatrial node HCN channel blockade is reachable: it is whether the clinic considers Corlanor the right add-on, how the dispense is set up, and how the titration schedule fits into a UAE family's life.

This page explains how the pathway works in 2026 for a UAE-resident patient: who qualifies, where the prescribing cardiologist conversation happens, how Corlanor or Procoralan is dispensed, what the realistic out-of-pocket exposure band is in AED, what to monitor on therapy (heart rate at every titration step, the distinctive luminous phosphene visual phenomenon, atrial fibrillation surveillance), and how the indefinite oral treatment course settles into the family's heart-failure routine. It is concierge documentation written for a family already in conversation with a treating cardiologist who wants the operational reality laid out plainly.

Why Corlanor, and why now

Corlanor is ivabradine, a selective sinoatrial node HCN channel blocker. The HCN channel (hyperpolarization-activated cyclic nucleotide-gated channel) carries the funny current (I-f) that drives the spontaneous diastolic depolarisation of the sinoatrial pacemaker cells. Ivabradine binds and blocks this channel selectively. The clinical translation is a clean reduction in resting and exercise heart rate, with no negative inotropic effect, no effect on cardiac conduction below the SA node, and no effect on blood pressure. That selectivity is what distinguishes ivabradine from beta-blockers: a heart-failure patient already on a maximally tolerated beta-blocker dose whose resting heart rate remains at 70 bpm or above has a real residual prognostic burden that ivabradine specifically addresses without compounding the negative inotropic load.

The FDA approved Corlanor in April 2015 for chronic HFrEF (NYHA II to IV, stable, sinus rhythm, resting HR at or above 70 bpm, on maximally tolerated beta-blocker or with documented contraindication). The paediatric expansion to children at or above 6 months with stable symptomatic dilated cardiomyopathy heart failure followed in April 2019, with an oral solution formulation. The European Procoralan label (Servier) is similar in shape and is registered across several Gulf markets. UAE EDE registration is in place for the indication; Procoralan may be the brand name actually stocked at some UAE pharmacies.

The pivotal trial is SHIFT, which randomised 6,558 adults with HFrEF, NYHA II-IV, LVEF at or below 35%, sinus rhythm, resting HR at or above 70 bpm, on stable GDMT including beta-blocker, to ivabradine or placebo. The primary composite endpoint of cardiovascular death or heart-failure hospitalisation was significantly reduced.

Reserve Meds does not promote one heart failure agent over another. The page describes the Corlanor pathway because Corlanor is the medication the family has asked about.

What Corlanor is, in plain language

Corlanor for adults is an oral tablet. 5 mg starting dose, twice daily, with food, approximately twelve hours apart. The titration schedule is heart-rate-driven: at the 2-week review the resting HR is checked, and the dose is adjusted up to 7.5 mg BID if resting HR is still above 60 bpm, held at 5 mg BID if HR is in the 50 to 60 bpm window, reduced to 2.5 mg BID if HR has dropped to 50 to 60 bpm with bradycardia symptoms, or stopped if HR has fallen below 50 bpm. The maintenance dose range is 2.5 to 7.5 mg BID.

The paediatric oral solution formulation (children at or above 6 months) is weight-based, with a similar HR-driven titration philosophy adapted to paediatric HR norms. The prescribing paediatric cardiologist runs a tighter review cadence in the first months.

Treatment is indefinite. The patient who tolerates Corlanor and remains in HR target stays on it.

Eligibility at a UAE cardiologist clinic

For UAE-resident patients, the cardiology services apply the FDA label criteria with EDE local adaptation:

1. HFrEF with LVEF at or below 35% on recent echocardiogram. 2. NYHA II to IV, clinically stable. 3. Sinus rhythm. Atrial fibrillation as the predominant rhythm excludes ivabradine. 4. Resting heart rate at or above 70 bpm on maximally tolerated beta-blocker (bisoprolol, carvedilol, metoprolol succinate), or beta-blocker contraindication or intolerance. 5. GDMT optimised: ACE inhibitor / ARB / ARNI titrated, MRA added, SGLT2 inhibitor added where appropriate. 6. Baseline ECG without significant AV block or sinoatrial node dysfunction. 7. Pregnancy planning discussion for women of childbearing potential; effective contraception is required. 8. For paediatric patients: age at or above 6 months with stable symptomatic dilated cardiomyopathy heart failure on standard paediatric GDMT, on oral solution.

A UAE patient should arrive at the cardiology conversation with the most recent echocardiogram, ECG, NYHA statement, full medication list, NT-proBNP if measured, and any prior HF hospitalisation history.

The UAE prescribing and supply picture

In 2026 the UAE cardiology centres that prescribe ivabradine routinely include Cleveland Clinic Abu Dhabi Heart and Vascular Institute, SSMC, SKMC, Burjeel Medical City, Mediclinic City Hospital, American Hospital Dubai, and the DHA hospitals. Both Corlanor (Amgen) and Procoralan (Servier) are seen in UAE pharmacy stock; the brand actually dispensed depends on the pharmacy's import arrangement.

1. Prescribing physician: board-certified adult cardiologist for adult HFrEF, board-certified paediatric cardiologist for paediatric DCM. 2. Pharmacy dispensing: hospital pharmacy in inpatient or specialty outpatient setting; community pharmacy for ongoing refills. 3. Insurance pre-authorisation: Thiqa for Emirati nationals, Daman, and the major commercial insurers handle ivabradine as a cardiology-prescribed therapy. Turnaround is typically 7 to 14 days with a complete HFrEF pack. 4. Titration follow-up: cardiology at week 2, week 4, then every 3 months once on stable maintenance dose. ECG with HR check at every titration visit.

Cost band

Annual cash-pay cost for Corlanor or Procoralan in UAE private pharmacy is approximately USD 1,200 to 2,500 (approximately AED 4,400 to 9,200). This is much lower than specialty drugs in the Reserve Meds catalogue and Corlanor sits as a cardiology breadth entry rather than a high-cost specialty case. Where EDE-registered generic ivabradine is dispensed, the annual cost can drop a further 30 to 60 percent.

For Emirati nationals on Thiqa, the prescription is processed through the formulary. Commercial pre-authorisation is straightforward with a complete HFrEF pack.

What to expect on Corlanor

Heart-rate titration is the principal early-treatment task. The cardiologist sets the 5 mg BID starting dose, reviews at week 2, steps up to 7.5 mg BID if HR target is not reached. Most patients stabilise on 5 mg or 7.5 mg BID within 4 to 6 weeks.

The distinctive adverse event is the luminous phosphene phenomenon: bright spots or transient flashes in the visual field, more often peripheral, sometimes triggered by sudden light changes. This happens because the HCN channel ivabradine blocks in the SA node has a related isoform (I-h) in the retina. The phosphene is reported by approximately 14 percent of patients in pivotal trials, typically in the first 2 months, and most patients accommodate as the retina adjusts. Patients should be told to expect it, told it is not dangerous, and told to call the cardiologist only if the phenomenon is severe enough to affect driving or daily function.

Other monitoring axes: bradycardia (HR check at every visit), atrial fibrillation (small but real increased incidence; pulse self-checks and any palpitation flag warrants a call), and the rare hypertension signal worth a BP at every visit. In SHIFT the meaningful outcome was reduction in heart-failure hospitalisation and, in higher-HR subgroups, cardiovascular mortality.

When Corlanor is the wrong drug

For a UAE patient with resting HR below 70 bpm at baseline, with atrial fibrillation as the predominant rhythm, with severe hepatic impairment, with strong CYP3A inhibitors that cannot be stopped (clarithromycin, ketoconazole, itraconazole, ritonavir, nefazodone), or with second-or-third-degree AV block without pacemaker, the operational pathway shifts. Paediatric DCM patients should not start ivabradine until standard paediatric GDMT is optimised first.

For UAE patients where Corlanor is not the chosen add-on, the alternatives are continued GDMT optimisation, ARNI uptitration if not at target, SGLT2 inhibitor addition where not yet on board, or in advanced cases device therapy (CRT, ICD) or advanced heart-failure consultation. Reserve Meds does not push one heart-failure agent over another.

Closing CTA

We are a US-based concierge coordinator. We are not the prescriber and not the dispensing pharmacy. On a UAE Corlanor case we build the documentation pack with the treating cardiologist office (echocardiogram, ECG, NYHA statement, GDMT list, NT-proBNP, prior HF hospitalisation history), confirm the EDE dispensing pathway (Corlanor, Procoralan, or generic ivabradine as locally registered), run the insurance pre-authorisation conversation alongside the clinical pre-authorisation conversation, and stay with the case through the titration window and into stable maintenance. Clinical decisions remain with your treating cardiologist.

Reserve Meds's role

US-based concierge coordinator for cross-border specialty medicine. We are not the prescriber, not the dispensing pharmacy, and not the manufacturer. All clinical decisions remain with your treating physician.

Reserve Meds

reserved for you.

Composite case examples. This document is for general information only and does not constitute medical advice. Please consult your treating physician.

Reserve Meds is in pre-launch. Published timelines and cost ranges are indicative, not guarantees.
reservemeds.com · hello@reservemeds.com